ANTIGENIX AMERICA Inc.

PRODUCT INFORMATION SUMMARY

Anti-HLA-A,B,C

(Common	Framework	Region)	M300020	Purified 0	0.1 mg	
			M300030	FITC	100	Tests
			M300060	Biotin	100	Tests
			M300070	Phycoerythr	in 100	Tests

ANTIGEN DISTRIBUTION AND SPECIFICITY:

The HLA-A,B,C antigen is expressed on the surface of most nucleated cells. This monoclonal antibody recognizes a framework determinant residing on the heavy chain of the class I MHC antigen associated with beta 2-microglobulin.

CLONE: W6-32 Mouse IgG2a, kappa light chain
Derived from hybridization of murine myeloma (NS-1) cells with
spleen cells from BALB/c mice immunized with activated human tonsil
membrane. Affinity purified from murine ascites fluid.

CONJUGATION: Fluorescein isothiocyanate; Biotin ester, R-Phycoerythrin.

HANDLING AND STORAGE:

All forms are supplied as 1.0 mL of liquid. Fluorochromes should be protected from prolonged exposure to light. Reagents will be in a medium containing 0.01M phosphate-buffered saline, pH 7.4, 0.2% gelatin and 0.1% sodium azide. Dilute with buffer containing 0.1% BSA. All reagents in a liquid state should be stored at $2\text{--}8^\circ$ C when not in use.

PRODUCT USE:

- * For flow cytometry use 10-20 uL per test;
- * For immuno-histochemistry (reactive frozen sections only) , purified Anti-HLA-A,B,C should be diluted 1:80 (optimize) * ELISA and Immunoprecipitation

RESEARCH APPLICATIONS:

Use as a positive control in all immunologic procedures detecting cell surface antigens.

X-Reactivity: Non-Human Primates: Cyno Monkey, Chimp, Rhesus, Macaque, Baboon, Gorilla. Bovine, Shrew

FOR RESEARCH USE ONLY. NOT INTENDED FOR THERAPEUTIC OR DIAGNOSTIC USE.

www.antigenix.com